



Anti-Endothelial differentiation, sphingolipid G-protein-coupled receptor 5 (EDG5), Rabbit-Polyclonal Antibody

Catalog No. GB-30104

Antigen species: Human

Host species: Rabbit

Quantity: 100 μ g

Reactivity: Human

Form: Peptide affinity purified antibody

Applications: ELISA

Target description

This EDG5 protein is a member of the G protein-coupled receptors, as well as the EDG family of proteins. This protein participates in sphingosine 1-phosphate-induced cell proliferation, survival, and transcriptional activation.

Antigen

This polyclonal antibody was raised by immunizing rabbit with a synthetic peptide located on the putative extracellular domain of human EDG5.

Application

The antibody titer is more than 1000K for ELISA. It has not been tested in the other applications. However, for the first testing, we recommend 1/3,000 dilution for ELISA, 1/1,000 dilution for Western blot analysis (WB) of recombinant protein, 1/400 dilution for tissue extracts or cell lysates, 1/50 dilution for immunohistochemistry (IHC) staining on frozen cryosections, 1/30 dilution for IHC staining on paraffin embedded sections.

Related Products

1. Anti- endothelial differentiation, sphingolipid G-protein-coupled receptor, 1 (EDG1), pAb (GB-30083)
2. Anti- endothelial differentiation, sphingolipid G-protein-coupled receptor, 2 (EDG2), pAb (GB-30084)
3. Anti- endothelial differentiation, sphingolipid G-protein-coupled receptor, 3 (EDG3), pAb (GB-30085)

Ab dilution	Pre-bleed	Purified Ab
1:100	0.875	2.775
1:1,000	0.186	2.719
1:10,000	0.065	2.440
1:100,000	0.056	1.258
1:1,000,000	0.057	0.281
Titer		>1000K

Antigen is coated on EIA strips at 1 μ g per well.

ELISA Protocol

Add 200 μ l of blocking buffer and then wash wells with PBST buffer. Antiserum or peptide specific purified antibody GB-30104 is diluted in series as 10^2 ~ 10^6 folds and added in separate wells. Incubate antibody for 1hr. Wash unbound antibodies and add anti-rabbit IgG-HRP conjugate. Wash the plates and add substrate to develop color for 5 min. Read absorbance (ABS) at 650 nm. Amount of color is directly proportional to the amount of antibodies. Antibody is positive at >2 folds of ABS of control/Pre-Immune serum.

Storage

It is supplied peptide affinity purified antibody in lyophilized powder. Redissolve the powder with 100 microliter sterile water will restore to the original concentration 1mg/ml (1×PBS). Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

References

1. Takuwa,Y. Regulation of Rho family G proteins and cell motility by the Edg family of sphingosine 1-phosphate receptors. *Tanpakushitsu Kakusan Koso* 47 (4 Suppl): 496-502, 2002.
2. Spiegel, S. Sphingosine 1-phosphate: a ligand for the EDG-1 family of G-protein-coupled receptors. *Ann. N. Y. Acad. Sci.* 905: 54-60, 2000.

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE