



Anti-TGFB1-induced anti-apoptotic factor 1 (TIAF1), Chicken-Polyclonal Antibody

Catalog No. PY-10203 **Quantity:** 100µg **Applications tested:** Western blot
Antigen species: Human **Reactivity:** Human
Host species: Chicken **Form:** Antigen affinity purified antibody

Target description

This protein inhibits the cytotoxic effects of TNF-alpha and overexpressed TNF receptor adapters TRADD, FADD, and RIP. Involved in TGF-beta1 inhibition of IκBα expression and suppression of TNF-mediated IκBα degradation.

Antigen


This polyclonal antibody was raised by immunizing chicken with TIAF1 fusion protein (1-115 amino acids).

Application

Western blotting, tissue or cell immunostaining. Recommended starting dilution for Western blot analysis is 1:500, for tissue or cell staining is 1:200. Optimal working dilutions must be determined by the end user.

Related Products

1. Anti-Cytokeratin 8 (KRT8), Chicken pAb (PY-10149).
2. Anti-Cytokeratin 18 (KRT18), Chicken pAb (PY-10150).
3. Anti- Toll-like receptor 2 (TLR2), Chicken pAb (PY-10152)
4. Anti- Toll-like receptor 3 (TLR3), Chicken pAb (PY-10154)
5. Anti- Toll-like receptor 7 (TLR7), Chicken pAb (PY-10148)



E coli-derived fusion protein as test antigen.

Western blot Protocol

1. Block membrane with 5% non-fat milk in PBS-T for 1 hour at room temperature or longer at 4°C.
2. Incubate membrane with IgY antibodies at dilution of 1: 2,000 with 1% milk in PBS-T at R.T. for 1 h.
3. Rinse 3 times with PBS-T, then wash membrane with PBS-T, 5 min each, total of 3 times.
4. Incubate with 2nd antibody (goat-anti-IgY/Fc-HRP) at dilution 1:1,000 for ECL (with 1% milk PBS-T) at R.T. for 1h.
5. Rinse 3 times with PBS-T, then wash with PBS-T, 5 min each with shaking, total of 3 times.
6. Perform ECL detection of signal using Pierce ECL kit.

Storage

It is supplied as antigen affinity purified antibody in lyophilized powder. Redissolve the powder with 100 microliter sterile water will restore to the original concentration 1mg/ml (1×PBS). Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

References

Schultz, L., Khera, S., Sleeve, D., Heath, J. and Chang, N.S. TIAF1 and p53 functionally interact in mediating apoptosis and silencing of TIAF1 abolishes nuclear translocation of serine 15-phosphorylated p53 DNA Cell Biol. 23 (1), 67-74 (2004)